

Relativity and the Minimum Slope of the Isgur-Wise Function

B. D. Keister

Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213

(February 1, 2008)

Abstract

Sum rules based upon heavy quark effective theory indicate that the Isgur-Wise function $\xi(w)$ has a minimum slope $\rho_{\min}^2$ as $w \rightarrow 1$, where $\rho_{\min}^2 = 0$ for light degrees of freedom with zero spin and $\rho_{\min}^2 = \frac{1}{4}$ for light spin $\frac{1}{2}$. Quark-model studies reveal sources for a minimum slope from a variety of relativistic effects. In this paper the origins of the minimum slope in the sum-rule and quark-model approaches are compared by considering hadrons with arbitrary light spin. In both approaches the minimum slope increases with the light spin j_l , but there appears to be no detailed correspondence between the quark-model and sum-rule approaches.

Typeset using REVTeX

The physics of hadrons containing a single heavy quark displays additional symmetries beyond that of QCD [1]. Matrix elements of operators which act only on the heavy quark may depend upon the hadronic structure, but only trivially on the heavy quark mass in the limit that it is infinitely heavy. Transitions involving the heavy quark quantum numbers can be expressed in terms of universal functions $\xi(w)$, where $w = (v' \cdot v)$ and v and v' are the initial and final hadron four-velocities.

On very general grounds, in heavy quark effective theory (HQET) a sum rule can be derived [2,3] in which the Isgur-Wise function has the form $\xi(w) = 1 - \rho^2(w - 1) + \dots$ as $w \rightarrow 1$, and that the slope ρ^2 has a minimum value of $\frac{1}{4}$ for heavy quarks coupled to light degrees of freedom with spin $\frac{1}{2}$. Isgur and Wise [3] noted that this factor differs from the value zero obtained in nonrelativistic calculations, and also when the light spin $j_l = 0$, and they further speculated that it comes from Zitterbewegungen of the fermionic light degrees of freedom. The problem of a heavy quark coupled to light degrees of freedom with arbitrary spin was later addressed by Falk [4], who showed that for all light spins $j_l > 0$, both integral and non-integral, there is a non-vanishing minimum slope, thereby ruling out Zitterbewegungen as the relevant phenomenon.

The minimum slope of $\xi(w)$ has been investigated in the context of a quark model by Close and Wambach [23], who find $\rho = 1.19$ for their choice of model parameters. Le Yaouanc *et al.* [6–8] have examined systematically the relativistic contributions to the minimum slope using the framework of direct interactions [9,10]. They find three distinct sources: kinematics involving the bound wave function and the Jacobian for variable transformations, a normalization effect in the heavy quark current, and Wigner rotations of the light spectator. These authors used instant-form dynamics, and concentrated on systems where the light degrees of freedom have spin $\frac{1}{2}$.

In this paper we compare the origins of the minimum slope in the HQET sum-rule and quark-model approaches by considering systems where the light degrees of freedom have arbitrary spin. The only contribution to the minimum slope which depends upon the light spin j_l is the effect of Wigner rotations; its contribution to the slope grows with increasing j_l ,

a result also obtained in an analysis by Falk, who uses a Minkowski tensor representation of the light degrees of freedom, though the functional dependence upon j_l is different. However, it will also be seen that there are difficulties in making a detailed correspondence between the physics leading to the HQET sum-rule results and those of a model.

Our approach is similar to that of Le Yaouanc *et al.*, in that we use a Bakamjian-Thomas direct interaction [9], but we employ Dirac's point-form dynamics rather than the instant form. The advantage of this approach is that exposes the effects of non-infinite heavy masses in a transparent manner. In the heavy-quark limit, all forms of dynamics should yield the same results. The details of this approach are described elsewhere [11].

The plan of the paper is as follows: after a brief overview of the dynamical approach, a heuristic derivation is given of the contribution of Wigner rotations to the minimum slope for spin- $\frac{1}{2}$ light degrees of freedom; then the exact results for the minimum slope of Le Yaouanc *et al.* are extended to arbitrary spin and compared to those of the Minkowski tensor approach.

For calculational purposes, we employ non-covariant free-particle state vectors $|mj; \mathbf{v}\mu\rangle$ for a particle of mass m and spin j , which are normalized as follows:

$$\langle mj; \mathbf{v}'\mu' | mj; \mathbf{v}\mu \rangle = \delta_{\mu'\mu} \delta(\mathbf{v}' - \mathbf{v}). \quad (1)$$

We now consider matrix elements of the form $\langle M'j'; \mathbf{V}'\mu' | I^\alpha(0) | Mj; \mathbf{V}\mu \rangle$, where $I^\alpha(x)$ acts on the quantum numbers of the heavy quark.

In general, the heavy quark is bound to a light system with spin j_l and mass m_l . The matrix element is expressed in terms of single-particle states as follows:

$$\begin{aligned} \langle M'j'; \mathbf{V}'\mu' | I^\alpha(0) | Mj; \mathbf{V}\mu \rangle &= \int d\mathbf{v}'_h \int d\mathbf{v}_h \int d\mathbf{v}_l \langle M'j'; \mathbf{V}'\mu' | m'_h m_l \tfrac{1}{2} j_l; \mathbf{v}'_h \mathbf{v}_l \mu'_h \mu_l \rangle \\ &\quad \times \langle m'_h \tfrac{1}{2}; \mathbf{v}'_h \mu'_h | I^\alpha(0) | m_h \tfrac{1}{2}; \mathbf{v}_h \mu_h \rangle \langle m_h m_l \tfrac{1}{2} j_l; \mathbf{v}_h \mathbf{v}_l \mu_h \mu_l | Mj; \mathbf{V}\mu \rangle \end{aligned} \quad (2)$$

Here and henceforth repeated indices are summed. The non-interacting two-body mass of the system is

$$M_0^2 \equiv P_0^\alpha P_{0\alpha}; \quad P_0^\alpha \equiv m_h v_h^\alpha + m_l v_l^\alpha. \quad (3)$$

The zero subscript denotes a non-interacting system. The total velocity is

$$\mathbf{V} = \mathbf{P}/M_0. \quad (4)$$

The relative momentum $\mathbf{k}$ can be expressed in terms of $\mathbf{p}_l = m_l \mathbf{v}_l$ and a rotationless boost $L_c(\mathbf{V})$:

$$\begin{aligned} \mathbf{k} &= L_c^{-1}(\mathbf{V}) \mathbf{p}_l \\ &\equiv \mathbf{p}_l - \mathbf{V} \left[\omega_{m_l}(\mathbf{p}_l) - \frac{(\mathbf{V} \cdot \mathbf{p}_l)}{1 + V^0} \right], \end{aligned} \quad (5)$$

where

$$\omega_m(\mathbf{p}) \equiv \sqrt{m^2 + \mathbf{p}^2}. \quad (6)$$

In terms of $\mathbf{k}$, the non-interacting mass is

$$M_0 = \omega_{m_h}(\mathbf{k}) + \omega_{m_l}(\mathbf{k}). \quad (7)$$

The most general bound state wave function is expressed as follows:

$$\begin{aligned} \langle m_h m_l \tfrac{1}{2} j_l; \mathbf{v}_h \mathbf{v}_l \mu_h \mu_l | M j; \mathbf{V} \mu \rangle &= \left| \frac{\partial(\mathbf{V} \mathbf{k})}{\partial(\mathbf{v}_h \mathbf{v}_l)} \right|^{\frac{1}{2}} \delta(\mathbf{V} - \mathbf{V}(\mathbf{v}_h, \mathbf{v}_l)) \\ &\quad \times D_{\mu_h \bar{\mu}_h}^{\frac{1}{2} \dagger} [R_c(L_c(\mathbf{V}), \mathbf{k}/m_h)] D_{\mu_l \bar{\mu}_l}^{j_l \dagger} [R_c(L_c(\mathbf{V}), -\mathbf{k}/m_l)] \\ &\quad \times \langle \tfrac{1}{2} \bar{\mu}_h j_l \bar{\mu}_l | s \mu_s \rangle \langle l \mu_l s \mu_s | j \mu \rangle Y_{\mu_L}^L(\hat{\mathbf{k}}) \phi_L(k), \end{aligned} \quad (8)$$

where the Jacobian is

$$\left| \frac{\partial(\mathbf{v}_h \mathbf{v}_l)}{\partial(\mathbf{V} \mathbf{k})} \right| = \left(\frac{M_0^3}{m_h^3 m_l^3} \right) \frac{\omega_1(\mathbf{v}_h) \omega_1(\mathbf{v}_l)}{\omega_1(\mathbf{u}_h) \omega_1(\mathbf{u}_l) \omega_1(\mathbf{V})}; \quad \mathbf{u}_{h,l} \equiv \mathbf{k}/m_{h,l}, \quad (9)$$

and

$$\mathbf{V}(\mathbf{v}_h, \mathbf{v}_l) = \frac{1}{M_0} (m_h \mathbf{v}_h + m_l \mathbf{v}_l) \quad (10)$$

and $\phi_L(k)$ is a radial momentum wave function for an interacting state of mass M .

At this point we convert from LS to jj coupling and assume that the heavy quark is coupled to a “light spin” j_l which contains all angular momenta except for the spin of the heavy quark itself [12]. In that case we set $L = 0$ and Eq. 9 simplifies to

$$\begin{aligned}
\langle m_h m_l \frac{1}{2} j_l \mathbf{v}_h \mathbf{v}_l \mu_h \mu_l | M j; \mathbf{V} \mu \rangle &= \left| \frac{\partial(\mathbf{V} \mathbf{k})}{\partial(\mathbf{v}_h \mathbf{v}_l)} \right|^{\frac{1}{2}} \delta(\mathbf{V} - \mathbf{V}(\mathbf{v}_h, \mathbf{v}_l)) \\
&\times D_{\mu_h \bar{\mu}_h}^{\frac{1}{2} \dagger} [R_c(L_c(\mathbf{V}), \mathbf{k}/m_h)] D_{\mu_l \bar{\mu}_l}^{j_l \dagger} [R_c(L_c(\mathbf{V}), -\mathbf{k}/m_l)] \\
&\times \langle \frac{1}{2} \bar{\mu}_h j_l \bar{\mu}_l | s \mu_s \rangle (4\pi)^{-\frac{1}{2}} \phi_0(k),
\end{aligned} \tag{11}$$

The Wigner rotation has the following form:

$$R_c(L_c(\mathbf{v}), \mathbf{a}) = L_c^{-1}(\mathbf{a}') L_c(\mathbf{v}) L_c(\mathbf{a}); \quad a'^\mu = L_c^\mu{}_\nu(\mathbf{v}) a^\nu. \tag{12}$$

After some variable changes, we get

$$\begin{aligned}
\langle M' j'; \mathbf{V}' \mu' | I^\alpha(0) | M j; \mathbf{V} \mu \rangle &= (4\pi)^{-1} \int d\mathbf{k} \left| \frac{\partial \mathbf{V}'}{\partial \mathbf{v}'_h} \right|_{\mathbf{v}_l}^{-1} \left| \frac{\partial(\mathbf{v}_h \mathbf{v}_l)}{\partial(\mathbf{V} \mathbf{k})} \right|^{\frac{1}{2}} \left| \frac{\partial(\mathbf{V}' \mathbf{k}')}{\partial(\mathbf{v}'_h \mathbf{v}'_l)} \right|^{\frac{1}{2}} \\
&\times \langle \frac{1}{2} \bar{\mu}'_h j_l \bar{\mu}'_l | s' \mu'_s \rangle \phi_0^*(k') \\
&\times D_{\bar{\mu}'_h \mu'_h}^{\frac{1}{2}} [R_c(L_c(\mathbf{V}'), \mathbf{k}'/m'_h)] D_{\bar{\mu}'_l \mu'_l}^{j_l} [R_c(L_c(\mathbf{V}'), -\mathbf{k}'/m_l)] \\
&\times \langle m'_h \frac{1}{2}; \mathbf{v}'_h \mu'_h | I^\alpha(0) | m_h \frac{1}{2}; \mathbf{v}_h \mu_h \rangle \\
&\times D_{\mu_h \bar{\mu}_h}^{\frac{1}{2} \dagger} [R_c(L_c(\mathbf{V}), \mathbf{k}/m_h)] D_{\mu_l \bar{\mu}_l}^{j_l \dagger} [R_c(L_c(\mathbf{V}), -\mathbf{k}/m_l)] \\
&\times \langle \frac{1}{2} \bar{\mu}_h j_l \bar{\mu}_l | s \mu_s \rangle \phi_0(k),
\end{aligned} \tag{13}$$

where

$$\left| \frac{\partial \mathbf{V}'}{\partial \mathbf{v}'_h} \right|_{\mathbf{v}_l} = \left(\frac{m'_h}{M'_0} \right)^3 \left[1 - \frac{1}{v_h'^0} \mathbf{V}' \cdot (V'^0 \mathbf{v}_h' - v_h'^0 \mathbf{V}') \right]. \tag{14}$$

In the heavy quark limit, $m_h = m'_h = M_0 = M'_0 = M = M'$, and

$$\left| \frac{\partial \mathbf{V}'}{\partial \mathbf{v}'_h} \right|_{\mathbf{v}_l} \rightarrow 1; \quad \left| \frac{\partial(\mathbf{v}_h \mathbf{v}_l)}{\partial(\mathbf{V} \mathbf{k})} \right| \rightarrow \frac{1}{m_l^3} \frac{\omega_1(\mathbf{v}_l)}{\omega_1(\mathbf{u}_l)}, \tag{15}$$

with a similar result for the Jacobian involving primed variables. The remaining dependence of the integral upon quark masses occurs via the momentum $\mathbf{k}$ as defined in Eq. (5):

$$\begin{aligned}
\mathbf{k} &= \mathbf{p}_l - \mathbf{V} \left[\omega_{m_l}(\mathbf{p}_l) - \frac{(\mathbf{V} \cdot \mathbf{p}_l)}{1 + V^0} \right]; \\
\mathbf{k}' &= \mathbf{p}_l - \mathbf{V}' \left[\omega_{m_l}(\mathbf{p}_l) - \frac{(\mathbf{V}' \cdot \mathbf{p}_l)}{1 + V'^0} \right].
\end{aligned} \tag{16}$$

The momenta $\mathbf{k}$, $\mathbf{k}'$ and $\mathbf{p}_l$ are related to each other via Eq. (16). They depend in turn on the external velocities $\mathbf{V}$ and $\mathbf{V}'$, as well as m_l , *but they are independent of the heavy quark mass m_h* . The remaining terms in the Jacobians depend upon ratios of these three momenta to m_l , and thus they also do not depend upon m_h . The Wigner rotations of the heavy quark become the identity in the heavy-quark limit. Those of the light degrees of freedom depend upon $\mathbf{k}/m_l$, which may be non-negligible, as discussed below, but are independent of the heavy quark mass.

We now investigate the contributions from the model to $\xi(w)$ as $w \rightarrow 1$. Since $\xi(w)$ is independent of the heavy quark mass, it is sufficient to consider the case of elastic scattering from a heavy pseudoscalar meson. In the point form, all generators of the homogeneous Lorentz group are kinematic, so that any component of the current matrix element in any frame can be used to calculate invariant quantities. We therefore evaluate matrix elements of the charge operator in the Breit frame, where $\mathbf{V} = -\mathbf{V}' = \frac{1}{2}\mathbf{A} = \frac{1}{2}\mathbf{q}/M$, and $\mathbf{A}^2 = 2(w-1)$.

Before considering the general case of arbitrary light spin, we present here a heuristic derivation of the minimum slope for light spin $\frac{1}{2}$. With the normalization choice of Eq. (1), the Isgur-Wise function becomes, in the heavy quark limit,

$$\xi(w) = (4\pi)^{-1} \int d\mathbf{k} \phi_0^*(k') D_{\bar{\mu}_l' \mu_l}^{\frac{1}{2}} [R_c(L_c(\frac{1}{2}\mathbf{A}), -\mathbf{k}'/m_l)] D_{\mu_l \bar{\mu}_l}^{\frac{1}{2}\dagger} [R_c(L_c(-\frac{1}{2}\mathbf{A}), -\mathbf{k}/m_l)] \phi_0(k). \quad (17)$$

For definiteness, we use an S -wave harmonic oscillator wave function of the form $\phi_0(k) = \exp(-k^2/2\beta^2)/\sqrt{N}$. The use of an explicit wave function introduces model dependence, but for the moment we are only interested in a heuristic derivation of the minimum slope.

In the nonrelativistic limit, the slope of $\xi(w)$ as $w \rightarrow 1$ is an overall size effect due to the conventional form factor $F(\mathbf{q}^2) = 1 - \frac{1}{6}\mathbf{q}^2 R^2 + \dots$. In the harmonic oscillator model, this is

$$F(\mathbf{q}^2) = 1 - \frac{1}{4} \left(\frac{m_l}{M} \right)^2 \frac{\mathbf{q}^2}{\beta^2} + \dots, \quad (18)$$

which yields $[\mathbf{q}^2 = 2M^2(w-1)]$:

$$\xi_{NR}(w) = 1 - \frac{1}{2} \frac{m_l^2}{\beta^2} (w-1) + \dots. \quad (19)$$

While there are several contributions of relativistic origin to the slope of $\xi(w)$, we consider only that of Wigner rotations for this heuristic discussion. In the limit $w \rightarrow 1$, the Wigner rotation angle is, to lowest order in k/m_l ,

$$\theta_R \rightarrow \left| \frac{\mathbf{V} \times \mathbf{k}}{2m_l} \right|, \quad (20)$$

For the primed variables, the rotation axis is the same, and the rotation angle has the same magnitude and opposite sign, with the result that the product $D(\theta_R)D^\dagger(-\theta_R)$ can be replaced by a single $D(2\theta_R)$. The leading contribution in powers of $(w - 1)$ comes from the diagonal elements $D_{\pm\frac{1}{2}\pm\frac{1}{2}}^{\frac{1}{2}}(2\theta_R) = \cos \theta_R \rightarrow 1 - \frac{1}{2}\theta_R^2$. For an S -wave harmonic oscillator ground state, $\langle 0 | (\mathbf{V} \times \mathbf{k})^2 | 0 \rangle = \mathbf{V}^2 \beta^2$, and the net multiplicative contribution to $\xi(w)$ is

$$\xi_W(w) = 1 - \frac{1}{16} \frac{\beta^2}{m_l^2} (w - 1) + \dots \quad (21)$$

Combining the factors in Eqs. (19) and (21) yields

$$\xi(w) = \xi_W(w)\xi_{NR}(w) = 1 - \frac{1}{2} \frac{m_l^2}{\beta^2} (w - 1) - \frac{1}{16} \frac{\beta^2}{m_l^2} (w - 1) + \dots \quad (22)$$

Note that the slope receives contributions both from factors $\frac{m_l^2}{\beta^2}$ and $\frac{\beta^2}{m_l^2}$. In the extreme nonrelativistic limit, $\beta \rightarrow 0$, and the slope becomes unbounded due to the arbitrarily large geometric size proportional to $\frac{1}{\beta}$. There exists a sum rule which provides an upper bound on the slope [13–16], implying that this limit of the model is unphysical. In addition, the form factor contribution alone, which is present in a nonrelativistic calculation, provides no lower bound to the slope. The presence of the relativistic correction via the Wigner rotation means that there is a minimum slope of $\xi(w) = 1 - \rho^2(w - 1) + \dots$. Minimizing the coefficient of $(w - 1)$ in Eq. (22) gives

$$\rho^2 \geq \frac{\sqrt{2}}{4}, \quad (23)$$

which differs by a factor $\sqrt{2}$ from the original Bjorken sum rule [2,3]. However, the original sum rule receives additional QCD corrections [14–17] which raise the lower bound of the slope to approximately 0.4. Since a direct-interaction quark model involves effective degrees

of freedom which incorporate QCD corrections implicitly, a comparison of $\sqrt{2}/4 = 0.35$ to 0.4 can be taken as reasonable agreement, but little can be said beyond this of a quantitative nature. In any event, there are other contributions beyond those of this heuristic analysis.

We now turn to systems containing light degrees of freedom with higher spin. In this case there can be two or more distinct functions $\xi(w)$. For spin-one light spectators, this is well known [18–21]. The number n_ξ of independent Isgur-Wise functions can be obtained by counting the number of non-negative values of μ_l , i.e., the number of unique diagonal elements of the corresponding Wigner rotation. For spin-zero and spin-one mesons with $j_l = \frac{1}{2}$, and spin- $\frac{1}{2}$ baryons with $j_l = 0$, $n_\xi = 1$. For spin-one mesons with $j_l = \frac{3}{2}$, and spin- $\frac{1}{2}$ baryons with $j_l = 1$, $n_\xi = 2$. For now, we consider only the matrix element for fully aligned spins, i.e., $j_{\text{hadron}} = j_l + \frac{1}{2}$. The relevant Wigner rotation of $2\theta_R$ is [22]

$$D_{j_l j_l}^{j_l}(2\theta_R) = (\cos \theta_R)^{2j_l} \rightarrow 1 - j_l \theta_R^2. \quad (24)$$

Our heuristic minimization approach yields the following behavior for $\xi(w)$:

$$\xi_W(w) = 1 - \frac{\sqrt{j_l}}{2} \frac{\beta^2}{m_l^2} (w - 1) + \dots \quad (25)$$

For spins which are not fully aligned, the expression for the minimum slope becomes more complicated but is always less than $\sqrt{j_l}/2$.

We now consider arbitrary light spin with full relativistic effects. For spin- $\frac{1}{2}$ spectators, Le Yaouanc *et al.* find that a careful analysis yields the following result for $\rho_{\min}^2$:

$$\begin{aligned} \rho_{\min}^2 &= \rho_{\text{space}}^2 + \rho_{\text{Wigner}}^2 + \rho_{\text{quark}}^2 \\ &= \frac{1}{3} + \frac{1}{6} + \frac{1}{4}, \end{aligned} \quad (26)$$

where ρ_{space}^2 contains contributions due to boosts of wave function argument and Jacobians for variable transformations, ρ_{Wigner}^2 is the contribution from spectator Wigner rotations, and ρ_{quark}^2 is a normalization effect from the heavy quark current. They find that this minimum corresponds to arbitrarily large internal momenta ($\beta \rightarrow \infty$ in the harmonic oscillator picture). Given this result, one expects no change in the value of β which minimizes ρ^2 when

the contribution of the Wigner rotation to ρ^2 is altered by a change in its coefficient from $\frac{1}{2}$ to some other value of j_l . We thus expect that for general j_l and fully aligned spins,

$$\rho_{\min}^2 = \frac{1}{3} + \frac{j_l}{3} + \frac{1}{4}. \quad (27)$$

The slope grows with j_l for large j_l , in agreement with Falk. However, it is difficult to make a detailed comparison between the tensor/sum-rule approach and quark models of the sort described here. In the tensor/sum-rule approach the minimum slope vanishes when $j_l = 0$; this would correspond to $\rho_{\text{space}}^2 = \rho_{\text{quark}}^2 = 0$ in the quark model. In the nonrelativistic sense that ρ_{space}^2 represents a geometric radius, this limit corresponds to $\beta \rightarrow \infty$. But we have already seen that there are contributions to ρ_{space}^2 which are either independent of β , or which have a non-vanishing limit as $\beta \rightarrow \infty$. As noted by Le Yaouanc *et al.*, the additional non-vanishing contributions ρ_{space}^2 and ρ_{quark}^2 correspond to the *compositeness* of the hadron as described by the quark model. By contrast, the analyses based upon a tensor representation describe the hadron as a fully factorized product of heavy quark and spectator degrees of freedom. In a sense they correspond to the wave function at zero separation in the quark model, i.e., the limit of arbitrarily large internal momenta for which $\rho_{\text{Wigner}}^2 = \frac{1}{6}$. This reasoning may represent a partial connection between the quark model and the tensor/sum-rule approaches, but the difference between $\frac{1}{6}$ in the former and $\frac{1}{4}$ in the latter remains unexplained.

In summary, the effect of relativity attributed to the minimum slope of the Isgur-Wise function as obtained from sum rules can be seen from the viewpoint of a simple model to arise naturally from Wigner rotations of the light degrees of freedom. Direct-interaction models have other contributions to the minimum slope which are of relativistic origin, but they are independent of the light spin and thus have no counterpart in the HQET sum-rule treatment. It may well be that any composite model of a heavy hadron will yield a minimum slope of $\xi(w)$ which exceeds the sum-rule value.

The author wishes to thank N. Isgur for asking about the source of a minimum slope in models of the sort considered here, G. C. Boyd and W. Roberts for helpful comments, and

O. Pène for correspondence concerning Refs. [6–8]. This work was supported in part by the U.S. National Science Foundation under Grant PHY-9319641.

REFERENCES

- [1] N. Isgur and M. B. Wise, in *B Decays*, edited by S. Stone (World Scientific, Singapore, 1991), p. 158.
- [2] J. D. Bjorken, invited talk at Les Rencontres de Physique de la Vallée d'Aoste, La Thuile, Italy [SLAC Report No. SLAC-PUB-5278, 1990 (unpublished)]; J. D. Bjorken, I. Dunietz and J. Taron, Nucl. Phys. **B371**, 111 (1992).
- [3] N. Isgur and M. B. Wise, Phys. Rev. D **43**, 819 (1991).
- [4] A. F. Falk, Nucl. Phys. **B378**, 79 (1992).
- [5] F. E. Close and A. Wambach, Nucl. Phys. **B412**, 169 (1994).
- [6] A. Le Yaouanc, L. Oliver, O. Pène and J.-C. Raynal, Phys. Lett. **B365**, 319 (1995).
- [7] A. Le Yaouanc, L. Oliver, O. Pène and J.-C. Raynal, Phys. Lett. **B386**, 304 (1996).
- [8] V. Morénas, A. Le Yaouanc, L. Oliver, O. Pène and J.-C. Raynal, Phys. Lett. **B365**, 315 (1996).
- [9] B. Bakamjian and L. H. Thomas, Phys. Rev. **92**, 1300 (1953).
- [10] B. D. Keister and W. N. Polyzou, Adv. Nucl. Phys. **20**, 225 (1991).
- [11] B. D. Keister, Phys. Rev. D **46**, 3188 (1992).
- [12] D. Scora and N. Isgur, Phys. Rev. D **52**, 2783 (1995).
- [13] M. B. Voloshin, Phys. Rev. D **46**, 3062 (1992).
- [14] I. Bigi, M. Shifman, N. G. Uraltsev and A. Vainshtein, Phys. Rev. D **52**, 196 (1995).
- [15] A. G. Grozin and G. P. Korchemsky, Phys. Rev. D **53**, 1378 (1996).
- [16] C. G. Boyd, Z. Ligeti, I. Z. Rothstein and M. B. Wise, Phys. Rev. D **55**, 3027 (1997).
- [17] C. G. Boyd, B. Grinstein and A. V. Manohar, Phys. Rev. D **54**, 2081 (1996).

- [18] N. Isgur and M. B. Wise, Nucl. Phys. B **348**, 276 (1991).
- [19] H. Georgi, Nucl. Phys. B **348**, 293 (1991).
- [20] J. G. Körner, Nucl. Phys. **B** (Proc. Suppl.) **21**, 366 (1991)
- [21] T. Mannel, W. Roberts and Z. Ryzak, Nucl. Phys. B **355**, 38 (1991).
- [22] D. M. Brink and G. R. Satchler, *Angular Momentum, 2nd Edition* (Oxford University Press, London, 1968).
- [23] F. E. Close, *An Introduction to Quarks and Partons* (Academic Press, New York, 1979), and references therein.